

References Cited:

1. Discodermolide: A new bioactive polyhydroxylated lactone from the marine sponge *Discodermia dissolute*. Gunasekera, S. P.; Gunasekera, M.; Longley, R. E.; Schulte, G. K. *J. Org. Chem.* **1991**, *56*, 1346.
2. Discodermolide—a new, marine-derived immunosuppressive compound. I. In vitro studies. Longley, R. E.; Caddigan, D.; Harmody, D.; Gunasekera, M.; Gunasekera, S. P.; *Transplantation* **1991**, *52*, 650.
3. Discodermolide, a cytotoxic marine agent that stabilizes microtubules more potently than taxol. ter Haar, E.; Kowalski, R. J.; Hamel, E.; Lin, C. M.; Longley, R. E.; Gunasekera, S. P.; Rosenkranz, H. S.; Day, B. W. *Biochemistry* **1996**, *35*, 243.
4. The microtubule-stabilizing agent discodermolide competitively inhibits the binding of paclitaxel (Taxol) to tubulin polymers, enhances tubulin nucleation reactions more potently than paclitaxel, and inhibits the growth of paclitaxel-resistant cells. Kowalski, R. J.; Giannakakou, P.; Gunasekera, S. P.; Longley, R. E.; Day, B. W.; Hamel, E.; *Mol. Pharmacol.* **1997**, *52*, 613.
5. Chemical biology of the epothilones. Nicolaou, K. C.; Roshangar, F.; Vourloumis, D. *Angew. Chem. Int. Ed.*, **1998**, *37*, 2014.
6. Microtubule bundling. Sasse, F. *Current Biology*, **2000**, *10*, R469.
7. Taxol and discodermolide represent a synergistic drug combination in human carcinoma cell lines. Martello, L. A.; McDaid, H. M.; Regl, D. L.; Yang, C. H.; Ment, D. T.; Pettus, R. R.; Kaufman, M. D.; Arimoto, H.; Danishefsky, S. J.; Smith, A. B. III; Horwitz, S. B. *Clin. Cancer Res.* **2000**, *6*, 1978.
8. Discodermolide: Just another microtubule-stabilizing agent? No! A lesson in synergy. Giannakakou, P.; Fojo, T. *Clin. Cancer Res.* **2000**, *6*, 1613.
9. Emerging microtubule-stabilizing agents for cancer chemotherapy. Myles, D. C. *Annual Reports in Med. Chem.*, **2002**, *37*, 125.
10. The development of a practical total synthesis of discodermolide, a promising microtubule-stabilizing anticancer agent. Paterson, I.; Florence, G. J. *Eur. J. Org. Chem.*, **2003**, 2193.

11. Syntheses of discodermolides useful for investigating microtubule binding and stabilization. Nerenberg, J. B.; Hung, D. T.; Nerenberg, J. B.; Schreiber, S. L. *J. Am. Chem. Soc.* **1996**, *118*, 11054.
12. Acyclic diastereoselective synthesis using tartrate ester modified crotylboronates. Double asymmetric reactions with α -methyl chiral aldehydes and synthesis of the C (19)-C (29) segment of rifamycin S. Roush, W. R.; Palkowitz, A. D.; Ando, K. *J. Am. Chem. Soc.* **1990**, *112*, 6348.
13. Evolution of a gram-scale synthesis of (+)-discodermolide. Smith, A. B. III; Beauchamp, T. J.; LaMarche, M. J.; Kaufman, M. D.; Qiu, Y.; Arimoto, H.; Jones, D. R.; Kobayashi, K. *J. Am. Chem. Soc.* **2000**, *122*, 8654.
14. Gram-scale synthesis of (+)-discodermolide. Smith, A. B. III; Kaufman, M. D.; Beauchamp, T. J.; LaMarche, M. J.; Arimoto, H. *Org. Lett.* **1999**, *1*, 1823.
15. Benyl Trichloroacetimidate, a versatile reagent for acid-catalysed benzylation of hydroxy-groups. Iversen, T.; Bundle, D. R. *J. Chem. Soc., Chem. Commun.* **1981**, 1240.
16. Total synthesis of the antimicrotubule agent (+)-discodermolide using boron-mediated aldol reactions of chiral ketones. Paterson, I.; Florence, G. J.; Gerlach, K.; Scott, J. P. *Angew. Chem. Int. Ed.* **2000**, *39*, 377.
17. Towards the total synthesis of Phorboxazoles A and B: stereocontrolled synthesis of a C20-C32 subunit. Paterson, I.; Arnott, E. A. *Tetrahedron Lett.* **1998**, *39*, 7185.
18. Polyketide Synthesis using the boron-mediated, anti-aldol reactions of lactate-derived ketones: total synthesis of (-)-ACRL toxin IIIB. Paterson, I.; Wallace, D. J.; Cowden, C. J. *Synthesis* **1998**, 639.
19. Direct synthesis of Z-unsaturated esters. A useful modification of the horner-emmons olefination. Still, W. C.; Gennari, C. *Tetrahedron Lett.* **1983**, *24*, 4405.
20. Preparation and use of 1-iodoalkyl ylide. Chen, J.; Wang, T.; Zhao, K. *Tetrahedron Lett.* **1994**, *35*, 2827.
21. Synthesis of medium ring ethers. 5. the synthesis of (+)-laurencin. Burton, J. W.; Clark, J. S.; Derrer, S.; Stork, T. C.; Bendall, J. G.; Holmes, A. B. *J. Am. Chem. Soc.* **1997**, *119*, 7483.

22. A stereoselective synthesis of (Z)-1-iodo-1-alkenes. Stork, G.; Zhao, K. *Tetrahedron Lett.* **1989**, *30*, 2173.
23. Stereoselective synthesis of (Ξ)- and (E)-1,3-alkadienes from aldehydes using organotitanium and lithium reagents. Ikeda, Y.; Ukai, J.; Ikeda, N.; Yamamoto, H. *Tetrahedron* **1987**, *43*, 723.
24. Aldehyde-selective addition of alkynylchromium compounds prepared by reduction of alkynyl halides with chromium(II) reagent. Takai, K.; Kuroda, T.; Nakatsukasa, S.; Oshima, K.; Nozaki, H. *Tetrahedron Lett.* **1985**, *26*, 5585.
25. Synthetic studies towards halichondrins. Aicher, T. D.; Kishi, Y. *Tetrahedron Lett.* **1987**, *28*, 3463.
26. The Peterson olefination reaction. Ager, D. *Org. React.* **1990**, *38*, 1.
27. Enantio- and diastereoselective aldol reaction of achiral ethyl and methyl ketones with aldehydes: the use of enol diethylpinocampheylborinates. Paterson, I.; Goodman, J. M.; Lister, M. A.; Schumann, R. C.; McClure, C. K.; Norcross, R. D. *Tetrahedron*, **1990**, *46*, 4663.
28. Total synthesis of Tartrolon. Mulzer, J.; Berger, M. *J. Am Chem. Soc.* **1999**, *121*, 8393.
29. A practical improvement, enhancing the large-scale synthesis of (+)-discodermolide: A third-generation approach. Smith, A.B. III; Kaufman, M. D.; Beauchamp, T. J.; LaMarche, M. J.; Arimoto, H. *Org. Lett.* **2003**, *5*, 4405.
30. 1,6-Asymmetric induction in boron-mediated aldol reactions: application to a practical total synthesis of (+)-discodermolide. Paterson, I.; Delgado, O.; Florence, G. J.; Lyothier, I.; Scott, J. P.; Sereinig, N. S. *Org. Lett.* **2003**, *5*, 35.
31. Paterson, I.; Delgado, O.; Florence, G. J.; Lyothier, I.; Scott, J. P.; Sereinig, N. S. *Org. Lett.* **2003**, *5*, 35.
32. Formal synthesis of (+)-discodermolide. Francavilla, C.; Chen, W.; Kinder, F. R. Jr. *Org. Lett.*, **2003**, *5*, 1233.
33. Simplified discodermolide analogues: Synthesis and biological evaluation of 4-epi-7-dehydroxy-14, 16-didemethyl-(+)-discodermolides as microtubule-stabilizing agents. Choy, N.; Shin, Y.; Nguyen, P. Q.; Curran, D. P.;

- Balanchandran, R.; Madiraju, C.; Day, B. W. *J. Med. Chem.*, **2003**, *46*, 2846-2864.
34. Acetylated analogues of the microtubule-stabilizing agent discodermolide: Preparation and biological activity. Gunasekera, S. P.; Longley, R. E.; Isbrucker, R. A. *J. Nat. Prod.* **2001**, *64*, 171.
35. The relationship between Taxol and (+)-discodermolide: Synthetic analogs and modeling studies. Martello, L. A.; LaMarche, M. J.; He, L.; Beauchamp, T. J.; Smith, A. B. III; Horwitz, S. B. *Chem. Biol.* **2001**, *8*, 843.