

REFERENCES

- Altmann, K. H.; Imwinkelried, R.; Kesselring, R.; Rihs, G. 1',6'-Methano Carbocyclic Thymidine: Synthesis, X-Ray Crystal Structure, and Effect on Nucleic Acid Duplex Stability. *Tetrahedron Lett.* **1994**, *35*, 7625–7628.
- Altmann, K.-H.; Kesselring, R.; Francotte, E.; Rihs, G. 4',6'-Methano Carbocyclic Thymidine: A Conformationally Constrained Building Block for Oligonucleotides. *Tetrahedron Lett.* **1994**, *35*, 2331–2334.
- Altona, C.; Sundaralingam, M. Conformational Analysis of the Sugar Ring in Nucleosides and Nucleotides. A New Description Using the Concept of Pseudorotation. *J. Am. Chem. Soc.* **1972**, *94*, 8205–8212.
- Altona, C.; Sundaralingam, M. Conformational Analysis of the Sugar Ring in Nucleosides and Nucleotides. Improved Method for the Interpretation of Proton Magnetic Resonance Coupling Constants. *J. Am. Chem. Soc.* **1973**, *95*, 2333–2344.
- Bolli, M.; Leumann, C. Triple-Helix Formation of Oligodeoxynucleotides Containing [(3'S,5'R)-2'-Deoxy-3',5'-ethano- α -D-ribofuranosyl]nucleosides ("Bicyclo-deoxynucleosides"). *Angew. Chem. Int. Ed. Engl.* **1995**, *34*, 694–696.
- Ezzitouni, A.; Barchi, J. J., Jr.; Marquez, V. E. A Simple Approach to 1',1'a-Methano Carbocyclic Thymidine. *Chem. Commun.* **1995**, 1345–1346.
- Ezzitouni, A.; Marquez, V. E. Conformationally Locked Carbocyclic Nucleosides Built on a Bicyclo[3.1.0]hexane Template with a Fixed Southern Conformation. Synthesis and Antiviral Activity. *J. Chem. Soc., Perkin Trans. 1* **1997**, 1073–1078.
- Ezzitouni, A.; Russ, P.; Marquez, V. E. (1S,2R)-[(Benzyloxy)methyl]cyclopent-3-enol. A Versatile Synthone for the Preparation of 4',1'a-Methano- and 1',1'a-Methanocarbocyclic Nucleosides. *J. Org. Chem.* **1997**, *62*, 4870–4873.
- Jacobson, K. A.; Ji, X.-d.; Li, A.-H.; Melman, N.; Siddiqui, M. A.; Shin, K.-J.; Marquez, V. E.; Ravi, R. G. Methanocarpa Analogues of Purine Nucleosides as Potent and Selective Adenosine Receptor Agonists. *J. Med. Chem.* **2000**, *43*, 2196–2203.
- Kim, H. S.; Ravi, R. G.; Marquez, V. E.; Maddileti, S.; Wihlborg, A.-K.; Erlinge, D.; Malmsjö, M.; Boyer, J. T.; Harden, T. K.; Jacobson, K. A. Methanocarpa Modification of Uracil and Adenine Nucleotides: High Potency of Northern Ring Conformation at P2Y₁, P2Y₂, P2Y₄, and P2Y₁₁ but Not P2Y₆ Receptors. *J. Med. Chem.* **2002**, *45*, 208–218.
- Koshkin, A. A.; Nielsen, P.; Meldgaard, M.; Rajwanshi, V. K.; Singh, S. K.; Wengel, J. LNA (Locked Nucleic Acid): an RNA Mimic forming Exceedingly Stable LNA:LNA Duplexes. *J. Am. Chem. Soc.* **1998**, *120*, 13252–13253.
- Koshkin, A. A.; Rajwanshi, V. K.; Wengel, J. Novel Convenient Syntheses of LNA [2.2.1]Bicyclo Nucleosides. *Tetrahedron Lett.* **1998**, *39*, 4381–4384.
- Koshkin, A. A.; Singh, S. K.; Nielsen, P.; Rajwanshi, V. K.; Kumar, R.; Meldgaard, M.; Olsen, C. E.; Wengel, J. LNA (Locked Nucleic Acids): Synthesis of the Adenine, Cytosine, Guanine, 5-Methylcytosine, Thymine, and Uracil Bicyclonucleoside Monomers, Oligomerisation, and Unprecedented Nucleic Acid Recognition. *Tetrahedron* **1998**, *54*, 3607–3630.
- Kværnø, L.; Wengel, J. Investigation of Restricted Backbone Conformations as an Explanation for the Exceptional Thermal Stabilities of Duplexes Involving LNA (Locked Nucleic Acid): Synthesis and Evaluation of Abasic LNA. *Chem. Commun.* **1999**, 657–658.
- Levitt, M.; Warshel, A. Extreme Conformational Flexibility of the Furanose Ring in DNA and RNA. *J. Am. Chem. Soc.* **1978**, *100*, 2607–2613.
- Marquez, V. E.; Ezzitouni, A.; Russ, P.; Siddiqui, M. A.; Ford, H., Jr.; Feldman, R. J.; Mitsuya, H.; George, C.; Barchi, J. J., Jr. HIV-1 Reverse Transcriptase Can Discriminate between Two Conformationally Locked Carbocyclic AZT Triphosphate Analogues. *J. Am. Chem. Soc.* **1998**, *120*, 2780–2789.
- Marquez, V. E.; Siddiqui, M. A.; Ezzitouni, A.; Russ, P.; Wang, J.; Wagner, R. W.; Matteucci, M. D. Nucleosides with a Twist. Can Fixed Forms of Sugar Ring Pucker Influence Biological Activity in Nucleosides and Oligonucleotides? *J. Med. Chem.* **1996**, *39*, 3739–3747.
- Meldgaard, M.; Wengel, J. Bicyclic Nucleosides and Conformational Restriction of Oligonucleotides. *J. Chem. Soc., Perkin Trans. 1* **2000**, 3539–3554.

- Nielsen, K. E.; Singh, S. K.; Wengel, J.; Jacobsen, J. P. Solution Structure of and LNA Hybridized to DNA: NMR Study of the d(CT^LGCT^LT^LCT^LGC):d(GCAGAAGCAG) Duplex Containing Four Locked Nucleotides. *Bioconjugate Chem.* **2000**, *11*, 228–238.
- Obika, S.; Nanbu, D.; Hari, Y.; Morio, K.-i.; In, Y.; Ishida, T.; Imanishi, T. Synthesis of 2'-O,4'-C-Methyleneuridine and -cytidine. Novel Bicyclic Nucleosides having a Fixed C3'-endo Sugar Puckering. *Tetrahedron Lett.* **1997**, *38*, 8735–8738.
- Ravi, R. G.; Kim, H. S.; Servos, J.; Zimmermann, H.; Lee, K.; Maddileti, S.; Boyer, J. T.; Harden, T. K.; Jacobson, K. A. Adenine Nucleotide Analogues Locked in a Northern Methanocarba Conformation: Enhanced Stability and Potency as P2Y₁ Receptor Agonists. *J. Med. Chem.* **2002**, *45*, 2090–2100.
- Rodriguez, J. B.; Marquez, V. E.; Nicklaus, M. C.; Barchi, J. J., Jr. Synthesis of Cyclopropane-fused Dideoxycarbocyclic Nucleosides Structurally Related to Neplanocin C. *Tetrahedron Lett.* **1993**, *34*, 6233–6236.
- Rodriguez, J. B.; Marquez, V. E.; Nicklaus, M. C.; Mitsuya, H.; Barchi, J. J., Jr. Conformationally Locked Nucleoside Analogues. Synthesis of Dideoxycarbocyclic Nucleoside Analogues Structurally Related to Neplanocin C. *J. Med. Chem.* **1994**, *37*, 3389–3399.
- Shin, K. J.; Moon, H. R.; George, C.; Marquez, V. E. Construction of the Bicyclo[3.1.0]hexane Template of a Conformationally Locked Carbocyclic Adenoside via an Olefin Keto-Carbene Cycloaddition. *J. Org. Chem.* **2000**, *65*, 2172–2178.
- Singh, S. K.; Nielsen, P.; Koshkin, A. A.; Wengel, J. LNA (Locked Nucleic Acids): Synthesis and High-Affinity Nucleic Acid Recognition. *Chem. Commun.* **1998**, 455–456.
- Tarköy, M.; Bolli, M.; Schweizer, B.; Leumann, C. Nucleic Acid Analogues with Constraint Conformational Flexibility in the Sugar-Phosphate Backbone ('Bicyclo-DNA'). Part 1. Preparation of (3'S,5'R)-2'-Deoxy-3',5'-ethano- α -D-ribonucleosides ('Bicyclonucleosides'). *Helv. Chim. Acta* **1993**, *76*, 481–510.
- Tarköy, M.; Leumann, C. Synthesis and Pairing Properties of Decanucleotides from (3'S,5'R)-2'-Deoxy-3',5'-ethano- α -D-ribofuranosyladenine and -thymine. *Angew. Chem. Int. Ed. Engl.* **1993**, *32*, 1432–1434.
- Wahlestedt, C.; Salmi, P.; Good, L.; Kela, J.; Johnsson, T.; Hökfelt, T.; Broberger, C.; Porreca, F.; Lai, J.; Ren, K.; Ossipov, M.; Koshkin, A.; Jacobsen, N.; Skouv, J.; Oerum, H.; Jacobsen, M. H.; Wengel, J. Potent and Nontoxic Antisense Oligonucleotides Containing Locked Nucleic Acids. *Proc. Natl. Acad. Sci. USA* **2000**, *97*, 5633–5638.
- Wengel, J. Synthesis of 3'-C- and 4'-C- Branched Oligodeoxynucleotides and the Development of Locked Nucleic Acid (LNA). *Acc. Chem. Res.* **1999**, *32*, 301–310.
- Chem*730 Proteins and Nucleic Acids. Nucleic Acid Architecture.
<http://www.chembio.uoguelph.ca/educmat/chm730/h730.htm>. (accessed October 2003).